

ECDEU ASSESSMENT MANUAL FOR PSYCHOPHARMACOLOGY Revised, 1976

William Guy, Ph.D.

Biometric Laboratory
The George Washington University
Kensington, Maryland

DATE DUE

MY 9 85 129112			
23 Jan 1988			
	45230		Printed in USA

U.S. DEPARTMENT OF HEALTH, EDUCATION, A
Public Health Service
Alcohol, Drug Abuse, and Mental Health Administrat
National Institute of Mental Health
Psychopharmacology Research Branch
Division of Extramural Research Programs
5600 Fishers Lane
Rockville, Maryland 20852

STUDY	PATIENT	FORM	PERIOD	RATER	HOSPITAL
		117			
(1-6)	(7-9)	(10-12)	(13-15)	(16-17)	(79-80)
PATIENT'S NAME					
RATER					
DATE					

**ABNORMAL INVOLUNTARY
MOVEMENT SCALE
(AIMS)**

INSTRUCTIONS: Complete Examination Procedure (reverse side) before making ratings. Code: 0 = None
1 = Minimal, may be extreme normal
2 = Mild
3 = Moderate
4 = Severe

MOVEMENT RATINGS: Rate highest severity observed. Rate movements that occur upon activation one less than those observed spontaneously.

		(Circle One)	CARD 01 (18-19)
FACIAL AND ORAL MOVEMENTS:	1. Muscles of Facial Expression e.g., movements of forehead, eyebrows, periorbital area, cheeks; include frowning, blinking, smiling, grimacing	0 1 2 3 4	(20)
	2. Lips and Perioral Area e.g., puckering, pouting, smacking	0 1 2 3 4	(21)
	3. Jaw e.g., biting, clenching, chewing, mouth opening, lateral movement	0 1 2 3 4	(22)
	4. Tongue Rate only increase in movement both in and out of mouth, NOT inability to sustain movement	0 1 2 3 4	(23)
EXTREMITY MOVEMENTS:	5. Upper (arms, wrists, hands, fingers) Include choreic movements, (i.e., rapid, objectively purposeless, irregular, spontaneous), athetoid movements (i.e., slow, irregular, complex, serpentine). Do NOT include tremor (i.e., repetitive, regular, rhythmic)	0 1 2 3 4	(24)
	6. Lower (legs, knees, ankles, toes) e.g., lateral knee movement, foot tapping, heel dropping, foot squirring, inversion and eversion of foot	0 1 2 3 4	(25)
TRUNK MOVEMENTS:	7. Neck, shoulders, hips e.g., rocking, twisting, squirming, pelvic gyrations	0 1 2 3 4	(26)
GLOBAL JUDGMENTS:	8. Severity of abnormal movements	None, normal 0 Minimal 1 Mild 2 Moderate 3 Severe 4	(27)
	9. Incapacitation due to abnormal movements	None, normal 0 Minimal 1 Mild 2 Moderate 3 Severe 4	(28)
	10. Patient's awareness of abnormal movements Rate only patient's report	No awareness 0 Aware, no distress 1 Aware, mild distress 2 Aware, moderate distress 3 Aware, severe distress 4	(29)
DENTAL STATUS:	11. Current problems with teeth and/or dentures	No 0 Yes 1	(30)
	12. Does patient usually wear dentures?	No 0 Yes 1	(31)

EXAMINATION PROCEDURE

Either before or after completing the Examination Procedure observe the patient unobtrusively, at rest (e.g., in waiting room).

The chair to be used in this examination should be a hard, firm one without arms.

1. Ask patient whether there is anything in his/her mouth (i.e., gum, candy, etc.) and if there is, to remove it.
2. Ask patient about the current condition of his/her teeth. Ask patient if he/she wears dentures. Do teeth or dentures bother patient now?
3. Ask patient whether he/she notices any movements in mouth, face, hands, or feet. If yes, ask to describe and to what extent they currently bother patient or interfere with his/her activities
4. Have patient sit in chair with hands on knees, legs slightly apart, and feet flat on floor. (Look at entire body for movements while in this position).
5. Ask patient to sit with hands hanging unsupported. If male, between legs, if female and wearing a dress, hanging over knees. (Observe hands and other body areas.)
6. Ask patient to open mouth. (Observe tongue at rest within mouth.) Do this twice.
7. Ask patient to protrude tongue. (Observe abnormalities of tongue movement.) Do this twice.
- 8. Ask patient to tap thumb, with each finger, as rapidly as possible for 10–15 seconds; separately with right hand, then with left hand. (Observe facial and leg movements.)
9. Flex and extend patient's left and right arms (one at a time.) (Note any rigidity and rate on DOTES.)
10. Ask patient to stand up. (Observe in profile. Observe all body areas again, hips included.)
- 11. Ask patient to extend both arms outstretched in front with palms down. (Observe trunk, legs, and mouth.)
- 12. Have patient walk a few paces, turn, and walk back to chair. (Observe hands and gait.) Do this twice.
- Activated movements