

Potential side effects associated with fluvoxamine (Luvox®)

| Common side effects* | Drug-placebo difference | Percentage of patients on fluvoxamine who reported it | Percentage of patients on placebo who reported it |
|--|-------------------------|---|---|
| Nausea | 26% | 40% | 14% |
| Somnolence (sleepiness) | 14% | 22% | 8% |
| Difficulty falling/staying asleep | 11% | 21% | 10% |
| Weakness/lack of energy | 8% | 14% | 6% |
| Nervousness | 7% | 12% | 5% |
| Abnormal ejaculation | 7% | 8% | 1% |
| Dyspepsia (indigestion) | 5% | 10% | 5% |
| Other side effects* | | | |
| Sweating | 4% | 7% | 3% |
| Anorexia (decreased appetite) | 4% | 6% | 2% |
| Tremor | 4% | 5% | 1% |
| Vomiting | 3% | 5% | 2% |
| Taste perversion | 2% | 3% | 1% |
| Tooth disorder (toothache, tooth extraction and abscess, and caries) | 2% | 3% | 1% |
| Vasodilatation | 2% | 3% | 1% |
| Difficulty achieving orgasm | 2% | 2% | 0% |
| Yawn | 2% | 2% | 0% |

| | | | |
|------------------------------------|-----------|----|----|
| Agitation | 1% | 2% | 1% |
| Chills | 1% | 2% | 1% |
| CNS stimulation | 1% | 2% | 1% |
| Decreased libido | 1% | 2% | 1% |
| Depression | 1% | 2% | 1% |
| Difficulty swallowing | 1% | 2% | 1% |
| Difficulty breathing | 1% | 2% | 1% |
| Hypertonia (increased muscle tone) | 1% | 2% | 1% |
| Impotence | 1% | 2% | 1% |
| Urinary Retention | 1% | 1% | 0% |

Adverse events that occurred during treatment with fluvoxamine are considered here to be side effects if they occurred at least twice as often as on placebo.

* Occurred in at least 5% more patients on fluvoxamine than in those on placebo.

** Occurred in at least 1% more patients on fluvoxamine than in those on placebo.

For important information including about potentially serious adverse effects of this medication, please see <http://www.medlineplus.gov/>.