

Potential side effects associated with lurasidone (brand name Latuda®)

(20 to 60 mg taken by itself to treat bipolar depression in adults)

Common side effects*	Drug-placebo difference	Percentage of patients on lurasidone who reported it	Percentage of patients on placebo who reported it
Akathisia (Restlessness or urge to move around)	6%	8%	2%
Other side effects*			
Diarrhea	3%	5%	2%
Extrapyramidal symptoms (Difficulty moving or slow movements)	3%	5%	2%
Anxiety	3%	4%	1%
Nasopharyngitis (Inflammation of nose and throat)	3%	4%	1%
Back pain	>2%	3%	<1%
Urinary tract infection	>1%	2%	<1%

*Adverse (undesirable) events that occurred during treatment with lurasidone are considered here to be side effects if they occurred *at least* twice as often as on placebo. A side effect is considered here to be “common” if the difference between how often it occurred on the medication versus on placebo is at least 5%. “Other” side effects included in this handout are those for which the difference between how often it occurred on the drug and on placebo is at least 1%. Side effects that occurred in less than 1% of patients, including potential serious side effects, are not covered by this handout.

For important information, including about potentially serious side effects of this medication, please make sure to also review the information at:

<https://medlineplus.gov/druginfo/meds/a611016.html>.

Disclaimer: The information in this handout is provided for use by medical professionals as only one tool in counseling their patients. It is not intended to be a complete list of possible side effects of this medication. Prescribing clinicians must consult the full Prescribing Information for this medication for its other side effects, including potentially serious ones.